

# Inhibition by the protein ceruloplasmin of lipid peroxidation stimulated by an $\text{Fe}^{3+}$ -ADP-adriamycin complex

Hiroko Nakano, Kouji Ogita, John M.C. Gutteridge\* and Minoru Nakano

*College of Medical Care and Technology, Gunma University, Maebashi, Gunma, Japan and \*National Institute for Biological Standards and Control, Holly Hill, Hampstead, London NW3 6RB, England*

Received 20 October 1983; revised version received 5 December 1983

Self-reduction of an  $\text{Fe}^{3+}$ -ADP-adriamycin complex under anaerobic conditions and reduction of ferricytochrome *c* by the complex under aerobic conditions were strongly inhibited by ceruloplasmin, but not by superoxide dismutase or albumin at the same protein concentration. Ceruloplasmin, a protein with ferroxidase activity, is able to catalyse oxidation of  $\text{Fe}^{2+}$  to the ferric state. The inhibitory activity of ceruloplasmin towards reactions stimulated by the complex suggests that  $\text{Fe}^{2+}$  is formed during the self-reduction process. As expected, the  $\text{Fe}^{3+}$ -ADP-adriamycin complex stimulated lipid peroxidation in which the  $\text{Fe}^{2+}$  moiety was implicated. This stimulation was again effectively prevented by ceruloplasmin but not by superoxide dismutase.

*Ceruloplasmin*

*Superoxide dismutase*  
*Lipid peroxidation*

*Adriamycin*  
*Oxygen radical*

*Ferroxidase activity*

## 1. INTRODUCTION

Ceruloplasmin, the copper-containing protein of extracellular fluids, has been shown to have important antioxidant properties towards peroxidising lipids [1-3]. These activities can be mainly ascribed to its known ferroxidase activity resulting in a catalysed oxidation of  $\text{Fe}^{2+}$  to the ferric state as well as an ability to inactivate copper ions [4].

The anthracycline antitumour antibiotic adriamycin has been shown, *in vivo*, to induce peroxidation of cardiac lipids [5]. Studies *in vitro*, however, have shown that adriamycin itself does not stimulate lipid peroxidation until it is first coordinated with  $\text{Fe}^{3+}$  [6]. Authors in [6,7] proposed a mechanism for adriamycin-induced lipid peroxidation involving an  $\text{Fe}^{3+}$ -chelated adriamycin complex in which the  $\text{Fe}^{3+}$  moiety of the complex could undergo self-reduction or enzymic reduction [6,7].

If ceruloplasmin can catalyse oxidation of the  $\text{Fe}^{2+}$  moiety in the chelated complex, then

adriamycin-iron-stimulated lipid peroxidation should be strongly inhibited by its ferroxidase activity. This work was undertaken to examine this proposal.

## 2. MATERIALS AND METHODS

### 2.1. Reagents

Superoxide dismutase (spec. act. 2900 units/mg), albumin (bovine, fatty acid-free), cytochrome *c* (horse heart, type VI) and ceruloplasmin (human, type III) were obtained from Sigma. Adriamycin (doxorubicin hydrochloride) was a gift from Kyowa Hakko Co. Ferrocyclochrome *c* was prepared by dithionite reduction of ferricytochrome *c* and passed through a Sephadex G-25 column ( $1.5 \times 12$  cm) which had been equilibrated with 10 mM Tris-HCl buffer at pH 7.5, using the same buffer to remove excess dithionite. During and after elution of ferrocyclochrome *c*, the buffer and eluate were gassed with nitrogen.

## 2.2. Phospholipid

Microsomal phospholipid was extracted from rat liver microsomes using chloroform-methanol [8] and was freed from contaminating lipids (free fatty acids, cholesterol and its ester) by a silicic acid column chromatography [9].

## 2.3. Incubation experiments

Components of the reaction mixture, such as phospholipid (as liposomes) and the  $\text{Fe}^{3+}$ -ADP complex, were prepared just before use as in [7].

Components of the co-ordination complex,  $\text{Fe}^{3+}$ -ADP or  $\text{Fe}^{3+}$ -ADP-adriamycin were expressed in molar concentrations. The  $\text{Fe}^{3+}$ -ADP-adriamycin complex prepared under our experimental conditions had a co-ordination ratio of about 2 for the adriamycin (ligand) to  $\text{Fe}^{3+}$  (co-ordination metal) [6]. A Thunberg-type cell with a sidearm was employed for anaerobic conditions. The contents (0.2 mM adriamycin and enzyme - when added - in 0.1 M Tris-HCl buffer, pH 7.5) of the cell were rapidly frozen using solid  $\text{CO}_2$ , followed by evacuation at 8 mm Hg at room temperature, until the contents were completely thawed. This was repeated 8 times in all. The reaction was initiated by mixing the  $\text{Fe}^{3+}$ -ADP complex held in the sidearm with the cell contents.

For all incubation experiments, the reaction was initiated by the addition of  $\text{Fe}^{3+}$ -ADP complex and continued at 37°C for the times indicated.

## 2.4. Assays

Self-reduction of the  $\text{Fe}^{3+}$ -ADP-adriamycin complex under anaerobic conditions was measured by the decrease in absorbance at 602 nm [6,7].

Oxygen consumption was measured in an 'Instech' oxygenometer (Model 102) with a Clark-type electrode [7].

Oxidation and reduction of cytochrome *c* were monitored by changes in absorbance at 550 nm [7]. Fatty acid composition was determined by a modification of the method in [10] using a Shimadzu gas chromatograph (Model GC-4CM).

## 3. RESULTS

### 3.1. Inhibition of self-reduction of the $\text{Fe}^{3+}$ -ADP-adriamycin complex by ceruloplasmin

When 0.2 mM adriamycin in 0.1 M Tris-HCl buf-

fer (pH 7.5) was mixed with the  $\text{Fe}^{3+}$  (0.1 mM)-ADP (1.67 mM) complex under both anaerobic and aerobic conditions it showed a characteristic absorption band at 602 nm [7]. Under anaerobic conditions the intensity of absorbance then decreased at 602 nm with increasing time, suggesting the self-reduction of the  $\text{Fe}^{3+}$ -ADP-adriamycin complex in which  $\text{Fe}^{3+}$  was converted to  $\text{Fe}^{2+}$  (fig.1, curve 1). Superoxide dismutase (0.05 mg/ml,  $1.6 \times 10^{-6}$  M) did not alter the original decay curve of the  $\text{Fe}^{3+}$ -ADP-adriamycin complex (fig.1, curve 2). The self-reduction process was biphasic, showing an initial slow reaction followed by a faster phase. The first response may represent the initiation of intramolecular electron transfer resulting in reduction of the iron moiety. Ceruloplasmin (0.05 mg/ml,  $3.3 \times 10^{-7}$  M) caused almost complete inhibition of the initial phase (fig.1, curve 3). However, inhibition did not last long, probably because of loss of ferroxidase activity under anaerobic conditions [11]. A further consideration for the reaction response is the stoichiometry between  $\text{Fe}^{2+}$  in the complex and copper ions in the enzyme. Ceruloplasmin (with 6-7 copper




Fig.1. Decrease of a characteristic absorption band (at 602 nm) for the  $\text{Fe}^{3+}$ -ADP-adriamycin complex under anaerobic conditions. The reaction mixtures and experimental conditions are described in section 2. 1,  $\text{Fe}^{3+}$ -adriamycin complex; 2,  $\text{Fe}^{3+}$ -ADP-adriamycin complex plus superoxide dismutase; 3,  $\text{Fe}^{3+}$ -ADP-adriamycin complex plus ceruloplasmin. Arrow indicates the time at which air was bubbled into the reaction mixture.

atoms/molecule) would, in this reaction, show a copper concentration corresponding to about 1/13 of the iron present. Little or no inhibition of the decrease in absorbance at 602 nm was observed with heat-denatured ceruloplasmin (not shown). In all experiments, intensity at  $A_{602\text{nm}}$  returned to the original level following re-oxygenation (fig.1).

### 3.2. Inhibition by ceruloplasmin of cytochrome *c* reduction by the $\text{Fe}^{3+}$ -ADP-adriamycin complex

Ferricytochrome *c* was reduced to ferrocyclochrome *c* by the  $\text{Fe}^{3+}$ -ADP-adriamycin complex under both anaerobic and aerobic conditions, suggesting an electron transfer from the  $\text{Fe}^{2+}$ -ADP-adriamycin complex (or a structurally related complex) produced by its self-reduction [7]. If oxidation of the  $\text{Fe}^{2+}$ -ADP-adriamycin complex is catalysed by ceruloplasmin, the reduction of




Fig.2. Effect of various proteins on the autooxidation of ferrocyclochrome *c* (A), and on cytochrome *c* reduction by the  $\text{Fe}^{3+}$ -ADP-complex (B), both in air. (A) The reaction mixture contained 0.018 mM ferrocyclochrome *c* (freshly prepared) and protein at 0.05 mg/ml (or none) in 0.1 M Tris-HCl buffer (pH 7.5). 1, ferrocyclochrome *c* (basal), basal plus superoxide dismutase or basal plus albumin; 2, basal plus ceruloplasmin; 3, basal plus heat-denatured ceruloplasmin. (B) The reaction mixture contained  $\text{Fe}^{3+}$  (0.1 mM)-ADP (1.67 mM)-adriamycin (0.2 mM), 0.018 mM ferricytochrome *c* and protein at 0.05 mg/ml (or none) in 0.1 M Tris-HCl buffer (pH 7.5). 1, ferricytochrome *c* plus  $\text{Fe}^{3+}$ -ADP-adriamycin complex (basal); 2, basal plus superoxide dismutase; 3, basal plus ceruloplasmin; 4, basal plus heat-denatured ceruloplasmin; 5, basal minus  $\text{Fe}^{3+}$ -ADP (control); 6, control plus ceruloplasmin.

cytochrome *c* by the  $\text{Fe}^{3+}$ -ADP-adriamycin complex should also be inhibited by ceruloplasmin. As shown in fig.2A, autooxidation of ferrocyclochrome *c* under aerobic conditions was promoted by native as well as heat-denatured ceruloplasmin. This may be attributed to non-specific activity associated with commercially obtained preparations of ceruloplasmin. As shown in fig.2B, intact ceruloplasmin caused a strong inhibition under aerobic conditions of the  $\text{Fe}^{3+}$ -ADP-adriamycin-induced cytochrome *c* reduction, while heat-denatured ceruloplasmin showed only a slight inhibitory effect. Little or no inhibition was observed following the addition of albumin or superoxide dismutase at the same protein concentration (0.05 mg/ml) as that for ceruloplasmin.

### 3.3. Inhibition by ceruloplasmin of $\text{Fe}^{3+}$ -ADP-adriamycin stimulated lipid peroxidation

Peroxidation of phospholipid liposomes in 0.1 M Tris-HCl buffer (pH 7.5) was monitored by an increase in oxygen consumption and changes in the fatty acid composition of the phospholipids. Oxygen consumption was markedly enhanced by




Fig.3. Oxygen consumption in the systems containing liposomes, the  $\text{Fe}^{3+}$ -ADP-adriamycin complex and protein (or none). (A) The reaction mixture contained liposomes (0.85  $\mu\text{mol/ml}$ ),  $\text{Fe}^{3+}$  (0.1 mM)-ADP (1.67 mM)-adriamycin (0.2 mM) complex and protein (0.05 mg/ml) or none in 0.1 M Tris-HCl buffer (pH 7.5). 1,  $\text{Fe}^{3+}$ -ADP-adriamycin and liposomes (basal) or basal plus albumin; 2, basal plus ceruloplasmin; 3, basal plus heat-denatured ceruloplasmin; 4, basal plus superoxide dismutase; 5, basal minus the  $\text{Fe}^{3+}$ -ADP-adriamycin complex; 6, basal minus liposomes. (B) The reaction mixture was the same as basal (1) in A. Arrow indicates the time at which ceruloplasmin (0.05 mg/ml) was added.

Table 1

Effect of ceruloplasmin on phospholipid fatty acid composition after peroxidation stimulated by an  $\text{Fe}^{3+}$ -ADP-adriamycin complex

Incubation system <sup>a</sup>	Fatty acid composition <sup>b</sup> (mol/100 mol palmitic acid)					
	16:0	18:0	18:1	18:2	20:4	22:6
S1 Suspension of phospholipid micelles	100	71	33	57	100	24
S2 Basal (after peroxidation)	100	72	33	51	67	12
S3 S2 + ceruloplasmin	100	72	33	55	84	18
S4 S2 + heat-denatured ceruloplasmin	100	70	34	54	75	14
S5 S2 + superoxide dismutase	100	73	33	53	76	16
S6 S2 + albumin	100	72	33	51	67	13

<sup>a</sup> The reaction mixtures and incubation conditions were the same as those described in the legend to fig.3. Samples were incubated for 15 min

<sup>b</sup> Results were expressed as means of 3 experiments which were reproducible to  $\pm 5\%$

the addition of the  $\text{Fe}^{3+}$ -ADP complex to the liposomal system containing 0.2 mM adriamycin. Oxygen consumption was substantially inhibited by the addition of ceruloplasmin (fig.3A) either before or during the reaction (fig.3B). It was also inhibited, but to a much smaller extent, by heat-denatured ceruloplasmin and native superoxide dismutase (fig.3A). Albumin, at the same protein concentration as ceruloplasmin, did not affect oxygen consumption. Under similar conditions, destruction of polyunsaturated fatty acids, especially arachidonic (20:4) and docosahexaenoic (22:6) acids, induced by the addition of  $\text{Fe}^{3+}$ -ADP-adriamycin was effectively prevented by ceruloplasmin but not by heat-denatured ceruloplasmin or intact superoxide dismutase (table 1).

#### 4. DISCUSSION

It has been reported previously [6,7] that an  $\text{Fe}^{3+}$ -ADP-adriamycin complex stimulates the peroxidation of phospholipid liposomes without the need to add a reducing system (e.g., NADH-dependent cytochrome P-450 reductase). Furthermore, it has been suggested that the co-ordination complex is aerobically converted to a perferryl ion-complex, via the  $\text{Fe}^{2+}$ -ADP-adriamycin radical, which stimulates the observed lipid peroxidation [7]. However, the  $\text{Fe}^{3+}$ -ADP-adriamycin complex which is formed in Tris-HCl buffer






can be dissociated into  $\text{Fe}^{3+}$ -ADP (or  $\text{Fe}^{3+}$ -ADP-phosphate) and adriamycin when present in a phosphate buffer. This has been observed to decrease stimulation of lipid peroxidation [12]. In addition, a simple  $\text{Fe}^{3+}$ -adriamycin complex has also been observed to stimulate lipid peroxidation [6,13].

Here, the strong inhibitory effect of ceruloplasmin towards the spontaneous decrease in absorbance at 602 nm (due to the self-reduction of the  $\text{Fe}^{3+}$ -ADP-adriamycin complex) under anaerobic conditions and also to reduction of cytochrome c by the  $\text{Fe}^{3+}$ -ADP-adriamycin complex under aerobic conditions has been demonstrated. This clearly indicates that the reduced iron moiety in the complex, which can be produced both by self-reduction and in the presence of an electron acceptor, is re-oxidised in a reaction catalysed by ceruloplasmin. Furthermore, ceruloplasmin offered a substantial protection against  $\text{Fe}^{3+}$ -ADP-adriamycin-stimulated lipid peroxidation as monitored by  $\text{O}_2$  uptake and estimation of changes in liposomal phospholipid fatty acids. This indicates that a species of  $\text{Fe}^{2+}$  (possibly the perferryl ion derived from  $\text{Fe}^{2+}$  co-ordinated with ADP and adriamycin) plays an important role in stimulating lipid peroxidation. In addition, a simple iron salt and iron chelate,  $\text{Fe}^{2+}$ -ADP, may also contribute.

The concentrations of superoxide dismutase used in the present experiments were far greater

than those required for its catalytic activity. Therefore, its inhibitory activity towards lipid peroxidation may not necessarily be of a truly enzymic nature since superoxide dismutase at  $5 \times 10^{-7}$  M had no effect on lipid peroxidation stimulated by the  $\text{Fe}^{3+}$ -ADP-adriamycin complex with [7] or without NADH-cytochrome P-450 reductase (not shown).

From the present results and those obtained previously [7] we can tentatively propose the following scheme:



where  $\text{O}_2^- \cdot \text{Fe}^{3+}\text{-ADP-ADM}^+$ , Cyt and E represent the perferryl ion complex, cytochrome *c* and ceruloplasmin, respectively. Reaction 5 should be considerably faster than reactions 2 and 4.

In conclusion, an iron-ADP-adriamycin complex can undergo a self-reduction to produce active

oxygen species able to stimulate lipid peroxidation. The reactivity of this species is dependent on the  $\text{Fe}^{2+}$  moiety which can be oxidised catalytically by the ferroxidase activity of the protein ceruloplasmin.

## REFERENCES

- [1] Stocks, J., Gutteridge, J.M.C., Sharp, R.J. and Dormandy, T.L. (1974) Clin. Sci. Mol., Med. 47, 223-233.
- [2] Gutteridge, J.M.C. (1977) Biochem. Biophys. Res. Commun. 77, 379-386.
- [3] Gutteridge, J.M.C., Richmond, R. and Halliwell, B. (1980) FEBS Lett. 112, 269-272.
- [4] Gutteridge, J.M.C. (1983) FEBS Lett. 157, 37-40.
- [5] Myers, C.E., McGuire, W.P., Liss, R., Ifram, I., Grotzinger, K. and Young, R. (1977) Science 197, 165-167.
- [6] Sugioka, K., Nakano, H., Noguchi, T., Tsuchiya, J. and Nakano, M. (1981) Biochem. Biophys. Res. Commun. 100, 1250-1258.
- [7] Sugioka, K. and Nakano, M. (1982) Biochim. Biophys. Acta 713, 333-343.
- [8] Folch, J., Lees, M.M. and Sloane-Stanley, G.H. (1957) J. Biol. Chem. 226, 497-509.
- [9] Hanahan, D.J., Dittner, J.C. and Warashima, E. (1957) J. Biol. Chem. 228, 685-700.
- [10] May, H.E. and McCay, P.B. (1968) J. Biol. Chem. 243, 2288-2295.
- [11] Osaki, S. (1966) J. Biol. Chem. 241, 5053-5059.
- [12] Sugioka, K., Nakano, H., Nakano, M., Tero-Kubota, S. and Ikegami, Y. (1983) Biochim. Biophys. Acta 753, 411-421.
- [13] Gutteridge, J.M.C. (1983) Biochem. Pharmacol. 32, 1949-1952.